

Information Leaflet



Liverpool Women's
NHS Foundation Trust

Early Medical Abortion Commenced at LWH (Up to 9 Weeks + 6 Days -Tablet Method)

PLEASE READ THIS LEAFLET CAREFULLY AND KEEP IT THROUGHOUT YOUR TREATMENT AS IT CONTAINS IMPORTANT INFORMATION REGARDING YOUR PROCEDURE AND AFTERCARE

POINTS TO BE AWARE OF BEFORE YOU START THE PROCEDURE

It is important before you take the first tablet that you are certain of your decision to proceed with the abortion.

There are known risks to the fetus from this medication and therefore, we must recommend that once you have taken the tablet(s) that you continue with the abortion.

It is recommended that you do not smoke or drink any alcohol once the procedure has commenced

Due to the unpredictability of the procedure, we recommend you inform us of any travel plans arranged.

If you are breastfeeding you may continue during the medical procedure as only small amounts of the drugs, Mifepristone and Misoprostol pass into the breast milk and are therefore not expected to cause adverse effects in the breastfed infant.

EARLY MEDICAL ABORTION PROCEDURE

First Part of your Treatment

You are advised to attend the Bedford Centre alone. If necessary- one support person may attend with you, however, you will be seen by the nurse alone. Children are not permitted on the department. You will be given a tablet called Mifepristone to swallow with

water whilst in the clinic. If you vomit within 30 minutes of taking it contact the clinic as you will need to return for another tablet.

Following this tablet you may experience some nausea and / or vomiting also bleeding and period type pain.

It is important to note that bleeding at this stage does not mean that you have passed the pregnancy therefore you should continue with the next stage of treatment as planned. If you have discomfort you can take over the counter pain relief such as Paracetamol and Ibuprofen as per package directions.

Final part of Treatment at home

You are advised to have a responsible adult stay with you at home when you take the Misoprostol tablets and to remain with you the rest of the day and overnight.

You will be given 4 tablets called Misoprostol to be taken at home 48 hours after the First medication and they can be taken in the following way

Option 1 - this is preferred method as you will experience fewer side effects. Place 4 of the Misoprostol tablets into the top of your vagina as high as you can place them. This can be done lying down or squatting. After two hours, repeat this process with the other two Misoprostol tablets.

Or

Option 2 - taken by mouth. Place 2 tablets on each side of your mouth (4 tablets in total), between cheek and gum and allow the tablets to dissolve for 30 minutes. If after 30 minutes they have not completely dissolved, swallow what is left with water. After two hours, repeat this process with the other two Misoprostol tablets.

What to expect after the final medication at home -

You will experience vaginal bleeding which can be heavy with blood clots and this may be accompanied by passage of the pregnancy tissue.

Period type (cramps) pain usually 1-2 hours after using the tablet. For any pain and discomfort, you experience you can use Paracetamol and Ibuprofen. Please read the patient information leaflet with the packet and take as directed.

You may experience some nausea, vomiting, dizziness, possible diarrhoea or cold or hot flushes. These are known side effects from these tablets and will not stop the process tablet from working.

For most people the procedure is likely to be completed within 4-6 hours of taking the Misoprostol, however it may take up to a few days.

We request that you contact the Bedford Centre for advice 7 days after taking the Misoprostol tablets if you have only had a slight blood loss without clots or you have not had any bleeding or you continue to have symptoms of pregnancy.

This may mean that the procedure has been unsuccessful.

KNOWN RISKS OF EARLY MEDICAL ABORTION

- The risk of a failed procedure and continuing pregnancy
- Excessive bleeding (haemorrhage) requiring blood transfusion,
- Retained pregnancy tissue requiring further treatment.
- Infection
- Undiagnosed ectopic pregnancy – this is rare
- Emotional / psychological distress following an abortion -If you are not coping, we would suggest that you contact your GP or Liverpool Bereavement Services on 0151 236 3932 so that counselling or support can be arranged.
- Risk of death is very rare.

We give you this information regarding the risks not to alarm you but we feel you would wish know so that you can make an informed decision. More detailed statistics are in the consent form you will have read and signed before starting the procedure.

FOLLOW UP INFORMATION

You will be given a pregnancy test which you should perform two weeks after you have taken the medication. It is very important that you follow this advice as this will determine whether the abortion has been successful.

Please only use the pregnancy test you are supplied with as they are different to the tests you can buy.

You must contact the Bedford Centre on **0151 708 9988** ext. **4509** if the pregnancy test is positive. We will arrange an appointment for you to be reviewed.

If you do not contact the Bedford Centre, we will assume that you have performed the home pregnancy test on the date requested and that it has shown a negative result

DISCHARGE ADVICE FOLLOWING EARLY MEDICAL ABORTION

1. As already discussed, you are advised to have a responsible adult, with you who is aware of the treatment and remain with you until the following morning.
2. You are advised not to travel long distances, especially by air, until you have performed a negative home pregnancy at 2 weeks post treatment. If you do decide to travel, we strongly recommend you inform your travel insurance provider that you are undergoing a procedure.
3. Bleeding following the procedure is very individual. It can last for up to 3 weeks as a continuation of the procedure, this is not a period. It is not unusual for the bleeding to stop and start during this time. With this bleeding we advise you to use sanitary towels not tampons.
4. You are advised to maintain your normal hygiene routine, however whilst you are bleeding when a showering it is important that you do not apply the jet of water directly to the vagina (douching) as this may increase the risk of infection.
5. You are advised to avoid sexual intercourse until you have a negative pregnancy test Using condoms may help to reduce the risk of infection.
6. You can get pregnant before your next period if adequate contraception is not used.

7. Your next period should occur 4 – 6 weeks following the procedure; however, this can be affected by the method of contraception you are using. The bleeding on this period may be heavier than you are normally used to.
8. It is advisable that you limit any strenuous activity including prolonged sports activities until your bleeding has settled.

If you should develop any of the following, If you should develop any of the following contact The **Bedford Clinic** on **0151-708-9988** extension **1130** between 08.30 and 16.30h. Outside these hours you must attend the Liverpool Women's Hospital **Gynaecology Emergency Department** or contact your local Accident & Emergency Department:

- Heavy and continuous bleeding that soaks through 2 or more pads in an hour for a period of 2 hours.
- Severe repeated or continuous abdominal pains - if not eased with the tablets you have taken for pain relief.
- Violent shivering attacks / chills
- High temperature
- Offensive (smelly) vaginal discharge

If you require any further advice regarding this method of abortion you can contact

The Bedford Centre on **0151-708-9988** extension **1130** between 08.30 and 16.30hrs Monday to Friday.

This leaflet can be formally translated on request via our Patient Experience Team, although response times to have information translated can vary. To request formal translation services or if you would like to make any suggestions or comments about the content of this leaflet, please contact the Patient Experience Team on 0151 702 4353 or by email at pals@lwh.nhs.uk

Liverpool Women's NHS Foundation Trust
Crown Street
Liverpool
L8 7SS

Tel: 0151 708 9988
Issue Date: 10/08/2022
Reference: Gyn_2024-363-V3
Review Date: 08/12/2026
© Liverpool Women's NHS Foundation Trust